


## What you'll learn about

- Product Rule in Integral Form
- Solving for the Unknown Integral
- Tabular Integration
- Inverse Trigonometric and Logarithmic Functions

## 6.3 Integration by Parts

### Product Rule in Integral Form

When  $u$  and  $v$  are differentiable functions of  $x$ , the Product Rule for differentiation tells us that

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$


Integrating both sides with respect to  $x$  and rearranging leads to the integral equation

$$\begin{aligned} \int \left( u \frac{dv}{dx} \right) dx &= \int \left( \frac{d}{dx}(uv) \right) dx - \int \left( v \frac{du}{dx} \right) dx \\ &= uv - \int \left( v \frac{du}{dx} \right) dx. \end{aligned}$$

When this equation is written in the simpler differential notation we obtain the following formula.



## Integration by Parts Formula

$$\int u \, dv = uv - \int v \, du$$

### EXAMPLE 1 Using Integration by Parts

Evaluate  $\int x \cos x \, dx$ .

#### SOLUTION

We use the formula  $\int u \, dv = uv - \int v \, du$  with

$$u = x, \quad dv = \cos x \, dx.$$

To complete the formula, we take the differential of  $u$  and find the simplest antiderivative of  $\cos x$ .

$$du = dx \quad v = \sin x$$

Then,

$$\int x \cos x \, dx = x \sin x - \int \sin x \, dx = x \sin x + \cos x + C.$$

*Now try Exercise 1.*

Q: is there an incorrect way to set this up?

A: Oh. Yes.

### EXPLORATION 1 Choosing the Right $u$ and $dv$

Not every choice of  $u$  and  $dv$  leads to success in antidifferentiation by parts. There is always a trade-off when we replace  $\int u dv$  with  $\int v du$ , and we gain nothing if  $\int v du$  is no easier to find than the integral we started with. Let us look at the other choices we might have made in Example 1 to find  $\int x \cos x dx$ .

1. Apply the parts formula to  $\int x \cos x dx$ , letting  $u = 1$  and  $dv = x \cos x dx$ . Analyze the result to explain why the choice of  $u = 1$  is never a good one.
2. Apply the parts formula to  $\int x \cos x dx$ , letting  $u = x \cos x$  and  $dv = dx$ . Analyze the result to explain why this is not a good choice for this integral.
3. Apply the parts formula to  $\int x \cos x dx$ , letting  $u = \cos x$  and  $dv = x dx$ . Analyze the result to explain why this is not a good choice for this integral.
4. What makes  $x$  a good choice for  $u$  and  $\cos x dx$  a good choice for  $dv$ ?

$$\begin{aligned} \textcircled{1} \quad u &= 1 \quad dv = x \cos x dx \\ du &= 0 \quad v = \textcircled{\text{---}} \end{aligned} \quad \left. \begin{aligned} \textcircled{2} \quad u &= x \cos x \quad dv = dx \\ du &= (1)(\cos x) dx + (-\sin x)(x) \\ v &= x \end{aligned} \right\}$$
$$\int u dv = uv - \int v du$$
$$= x^2 - \int x (\cos x) dx$$

In Exercises 1–10, find the indefinite integral.

1.  $\int x \sin x \, dx$   
 $-x \cos x + \sin x + C$

$$\int u \, dv = uv - \int v \, du$$

3.  $\int 3t e^{2t} \, dt$   $\frac{3}{2} t e^{2t} - \frac{3}{4} e^{2t} + C$  4

$$\begin{aligned} u &= x & dv &= \sin x \, dx \\ du &= dx & v &= -\cos x \\ &= -x \cos x - \int (-\cos x) \, dx \\ &= -x \cos x + \int \sin x \, dx \\ & & & + C \end{aligned}$$

$$3. \int 3t e^{2t} dt \quad \frac{3}{2} t e^{2t} - \frac{3}{4} e^{2t} + C$$

$$u = 3t \quad dv = e^{2t} dt$$
$$du = 3 dt \quad v = \frac{1}{2} e^{2t}$$

$$(3t) \left( \frac{1}{2} e^{2t} \right) - \int \frac{1}{2} e^{2t} (3) dt$$

$$\frac{3}{2} t e^{2t} - \frac{3}{4} e^{2t} + C$$

## EXAMPLE 2 Repeated Use of Integration by Parts

Evaluate  $\int x^2 e^x dx$ .

### SOLUTION

With  $u = x^2$ ,  $dv = e^x dx$ ,  $du = 2x dx$ , and  $v = e^x$ , we have

$$\int x^2 e^x dx = x^2 e^x - 2 \int x e^x dx.$$

The new integral is less complicated than the original because the exponent on  $x$  is reduced by one. To evaluate the integral on the right, we integrate by parts again with  $u = x$ ,  $dv = e^x dx$ . Then  $du = dx$ ,  $v = e^x$ , and

$$\int x e^x dx = x e^x - \int e^x dx = x e^x - e^x + C.$$

Hence,

$$\begin{aligned} \int x^2 e^x dx &= x^2 e^x - 2 \int x e^x dx \\ &= x^2 e^x - 2x e^x + 2e^x + C. \end{aligned}$$

$$5. \int x^2 \cos x \, dx$$

$$x^2 \sin x + 2x \cos x - 2 \sin x + C$$

5  $u = x^2 \quad dv = \cos x$   
 $du = 2x \, dx \quad v = \sin x$

$$7. \int 3x^2 e^{2x} \, dx$$

$$9. \int y \ln y \, dy \quad \frac{y^2}{2} \ln y - \frac{y^2}{4} + C$$

$$7. \frac{3}{2} x^2 e^{2x} - \frac{3}{2} x e^{2x} + \frac{3}{4} e^{2x} + C$$

$$\int u \, dv = uv - \int v \, du$$

$$\int x^2 \cos x \, dx = \underline{x^2 \sin x}$$

$$- \int 2x \sin x \, dx$$

$$u = 2x \quad dv = \sin x \, dx$$

$$du = 2 \quad v = -\cos x$$

$$\int 2x \sin x \, dx = \underline{2x(-\cos x)}$$

$$- \int -2 \cos x$$

$$+ 2 \sin x$$

$$= \underline{x^2 \sin x} + \underline{2 \cos x} - \underline{2 \sin x} + C$$



$$7. \int 3x^2 e^{2x} dx$$

derive...

anti

$$\frac{d}{dx} e^{2x} = 2 \cdot e^{2x}$$

<u>u</u>	<u>dv</u>
$3x^2$	$e^{2x}$
$6x$	$\frac{1}{2} e^{2x}$
$6$	$\frac{1}{4} e^{2x}$
$0$	$\frac{1}{8} e^{2x}$

$\frac{3}{2} x^2 e^{2x}$   
 $-\frac{3}{2} x e^{2x}$   
 $+\frac{3}{4} e^{2x}$

$$7. \frac{3}{2} x^2 e^{2x} - \frac{3}{2} x e^{2x} + \frac{3}{4} e^{2x} + C$$



$$6. \int x^2 e^{-x} dx$$

$$-x^2 e^{-x} - 2x e^{-x} - 2e^{-x} + C$$

$$\int u dv = uv - \int v du$$

$$u = \underline{x^2} \quad \underline{dv} = e^{-x}$$

$$du = 2x dx \quad v = -e^{-x}$$

$$= (x^2)(-e^{-x}) + \int 2x e^{-x} dx + C$$

$$u = \underline{2x} \quad \underline{dv} = e^{-x}$$

$$v = -e^{-x}$$

$$= -x^2 e^{-x} + 2x(-e^{-x}) - \int (-e^{-x})(2 dx)$$

$$= -x^2 e^{-x} - 2x e^{-x} + 2 \int e^{-x} dx$$

$$= -x^2 e^{-x} - 2x e^{-x} - \underline{2e^{-x}} + C$$

$$9. \int y \ln y \, dy = \frac{y^2}{2} \ln y - \frac{y^2}{4} + C$$

$u$	$dv$
$\ln y$	$y$
$-\frac{1}{x}$	$\frac{y^2}{2}$
$-\frac{1}{x^2}$	$\frac{y^3}{6}$

$u$	$dv$
$y$	$\ln y$

$$9. \int y \ln y \, dy \quad \frac{y^2}{2} \ln y - \frac{y^2}{4} + C$$

$$u = \ln y \quad dv = y$$
$$du = \frac{1}{y} dy \quad v = \frac{y^2}{2}$$

$$uv - \int v du$$

$$(\ln y) \left( \frac{y^2}{2} \right) - \int \frac{y^2}{2} \frac{1}{y} dy$$

$$\frac{y^2 \ln y}{2} - \int \frac{y}{2} dy$$

$$= \frac{y^2 \ln y}{2} - \frac{1}{2} \left( \frac{y^2}{2} \right) + C$$

### EXAMPLE 3 Solving an Initial Value Problem

Solve the differential equation  $dy/dx = x \ln(x)$  subject to the initial condition  $y = -1$  when  $x = 1$ . Confirm the solution graphically by showing that it conforms to the slope field.

#### SOLUTION

We find the antiderivative of  $x \ln(x)$  by using parts. It is usually a better idea to differentiate  $\ln(x)$  than to antidifferentiate it (do you see why?), so we let  $u = \ln(x)$  and  $dv = x dx$ .

$$\begin{aligned}y &= \int x \ln(x) dx \\&= \left(\frac{x^2}{2}\right) \ln(x) - \int \left(\frac{x^2}{2}\right) \left(\frac{1}{x}\right) dx \\&= \left(\frac{x^2}{2}\right) \ln(x) - \int \left(\frac{x}{2}\right) dx \\&= \left(\frac{x^2}{2}\right) \ln(x) - \frac{x^2}{4} + C\end{aligned}$$

Using the initial condition,

$$\begin{aligned}-1 &= \left(\frac{1}{2}\right) \ln(1) - \frac{1}{4} + C \\-\frac{3}{4} &= 0 + C \\C &= -\frac{3}{4}.\end{aligned}$$

Thus

$$y = \left(\frac{x^2}{2}\right) \ln(x) - \frac{x^2}{4} - \frac{3}{4}.$$

Figure 6.9 shows a graph of this function superimposed on a slope field for  $dy/dx = x \ln(x)$ , to which it conforms nicely.

**Now try Exercise 11.**

$y = -1, x = 1 \dots$

$$\int u dv = UV - \int v du$$

$$u = \ln t \quad dv = t^2$$
$$du = \frac{1}{t} dt \quad v = \frac{1}{3} t^3$$

$$\int (\ln t dt) (t^2) = \underline{(\ln t) \left(\frac{1}{3} t^3\right)} - \int \left(\frac{1}{3} t^3\right) \left(\frac{1}{t} dt\right)$$
$$= - \int \frac{1}{3} t^2 dt$$

$$10. \int t^2 \ln t dt \quad \underline{\frac{t^3}{3} \ln t} - \underline{\frac{t^3}{9}} + C$$

$$= - \underline{\frac{1}{9} t^3}$$

$$18. \int e^{-x} \cos x \, dx \quad \int u \, dv = uv - \int v \, du$$

$$u = e^{-x} \quad dv = \cos x \, dx$$

$$du = -e^{-x} \, dx \quad v = \sin x$$

$$\int (e^{-x})(\cos x \, dx) = (e^{-x})(\sin x) - \int (\sin x)(e^{-x} \, dx)$$

$$\underline{\sin x e^{-x}} = - \int -\sin x e^{-x}$$

$$\int u \, dv = uv - \int v \, du \quad \underline{\sin x e^{-x}} + \int \sin x e^{-x}$$

$$u = e^{-x} \, dx \quad dv = \sin x$$

$$du = -e^{-x} \quad v = -\cos x$$

$$\int (e^{-x})(\cos x \, dx) = \underline{(e^{-x})(-\cos x) - \int (-\cos x)(e^{-x} \, dx)}$$

$$\int e^{-x} \cos x \, dx = \sin x e^{-x} - (e^{-x})(\cos x) - \int (\cos x) e^{-x} \, dx$$

$$+ \int e^{-x} \cos x \, dx \quad + \int e^{-x} \cos x \, dx$$

$$2 \int e^{-x} \cos x \, dx = \frac{\sin x e^{-x}}{2} - \frac{\cos x e^{-x}}{2} + C$$

$$\frac{e^{-x}}{2} (\sin x - \cos x) + C$$

In Exercises 11–16, solve the initial value problem. Confirm your answer by checking that it conforms to the slope field of the differential equation.

11.  $\frac{dy}{dx} = (x + 2) \sin x$  and  $y = 2$  when  $x = 0$

$$y = \int (x+2) \sin x$$

$$\begin{aligned} u &= x+2 & \left\{ \begin{aligned} dv &= \sin x \\ du &= dx \end{aligned} \right. \\ v &= -\cos x \end{aligned}$$

13.  $\frac{du}{dx} = x \sec^2 x$  and  $u = 1$  when  $x = 0$

$$u v - \int v du$$

$$(x+2)(-\cos x) - \int (-\cos x) dx$$

$$y = -(x+2)(\cos x) + \sin x + C$$

15.  $\frac{dy}{dx} = x\sqrt{x-1}$  and  $y = 2$  when  $x = 1$

$$2 = -(2)(\cos 0) + \sin 0 + C$$

$$2 = -(2)(1) + 0 + C$$

$$C = 4$$

$$y = -(x+2)(\cos x) + \sin x + 4$$



**EXAMPLE 4 Solving for the Unknown Integral**Evaluate  $\int e^x \cos x \, dx$ .**SOLUTION**Let  $u = e^x$ ,  $dv = \cos x \, dx$ . Then  $du = e^x \, dx$ ,  $v = \sin x$ , and

$$\int e^x \cos x \, dx = e^x \sin x - \int e^x \sin x \, dx.$$

The second integral is like the first, except it has  $\sin x$  in place of  $\cos x$ . To evaluate it, we use integration by parts with

$$u = e^x, \quad dv = \sin x \, dx, \quad v = -\cos x, \quad du = e^x \, dx.$$

Then

$$\begin{aligned} \int e^x \cos x \, dx &= e^x \sin x - \left( -e^x \cos x - \int (-\cos x)(e^x \, dx) \right) \\ &= e^x \sin x + e^x \cos x - \int e^x \cos x \, dx. \end{aligned}$$

$$\int e^x \cos x \, dx$$

$$\int e^x \cos x \, dx = e^x \sin x + e^x \cos x - \int e^x \cos x \, dx$$

$$+ \int e^x \cos x \, dx$$

$$+ \int e^x \cos x \, dx$$

$$\underline{\underline{\underline{\int e^x \cos x \, dx = e^x \sin x + e^x \cos x}}}$$

$$19. \frac{e^x}{5} (2 \sin 2x + \cos 2x) + C$$

In Exercises 17–20, use parts and solve for the unknown integral.

$$17. \int e^x \sin x \, dx = \frac{e^x}{2} (\sin x - \cos x) + C$$

$$19. \int e^x \cos 2x \, dx =$$

$$u = e^x \quad dv = \cos 2x$$

$$du = e^x dx \quad v = \sin 2x \cdot \frac{1}{2}$$

$$\frac{1}{2} e^x \sin 2x - \int \frac{1}{2} \sin 2x \cdot e^x dx$$

$$u = e^x \quad dv = \sin 2x$$

$$du = e^x dx \quad v = -\cos 2x \cdot \frac{1}{2}$$

$$e^x (-\cos 2x) \frac{1}{2} - \int (-\cos 2x) \frac{1}{2} e^x dx$$

$$-\frac{1}{2} e^x \cos 2x + \frac{1}{2} \int \cos 2x e^x dx$$

In Exercises 17–20, use parts and solve for the unknown integral.

$$17. \int e^x \sin x \, dx = \frac{e^x}{2} (\sin x - \cos x) + C$$

$$18. \int e^{-x} \cos x \, dx = \frac{e^{-x}}{2} (\sin x - \cos x) + C$$

$$19. \int e^x \cos 2x \, dx$$

$$20. \int e^{-x} \sin 2x \, dx$$

(17)  $u = e^x \quad dv = \sin x \, dx$   
 $du = e^x \, dx \quad v = -\cos x$   
 $= (e^x)(-\cos x) - \int (-\cos x)(e^x) \, dx$   
 $= -e^x \cos x + \int \cos x e^x \, dx$   
 $du = e^x \, dx = \cos x \, dx$   
 $du = e^x \quad v = \sin x$

$\int e^x \sin x \, dx = -e^x \cos x + e^x \sin x$   
 $+ \int e^x \sin x \, dx$  "chasing the tail"  $\Rightarrow$   $-\int e^x \sin x \, dx$   


---

 $2 \int e^x \sin x \, dx = \frac{-e^x \cos x + e^x \sin x + C}{2}$   
 $\int e^x \sin x \, dx = \frac{-1}{2} \cos x + \frac{1}{2} e^x \sin x + C$

### EXAMPLE 5 Using Tabular Integration

Evaluate  $\int x^2 e^x dx$ .

#### SOLUTION

With  $u = x^2$  and  $dv = e^x$ , we list:

$u$ and its derivatives		$dv$ and its integrals
$x^2$	(+)	$e^x$
$2x$	(-)	$e^x$
$2$	(+)	$e^x$
$0$		$e^x$

Handwritten notes:  $u$  (with arrow pointing to  $x^2$ ),  $dv$  (with arrow pointing to  $e^x$ ), and a list of integrals:  $\sin x$ ,  $\cos x$ ,  $-\sin x$ ,  $-\cos x$  with arrows pointing to the  $e^x$  entries in the table.

We combine the products of the functions connected by the arrows according to the operation signs above the arrows to obtain

$$\int x^2 e^x dx = x^2 e^x - 2x e^x + 2e^x + C.$$

Compare this with the result in Example 2.

*Now try Exercise 21.*

### EXAMPLE 6 Using Tabular Integration

Evaluate  $\int x^3 \sin x \, dx$ .

#### SOLUTION

With  $f(x) = x^3$  and  $g(x) = \sin x$ , we list:

$f(x)$ and its derivatives		$g(x)$ and its integrals
$x^3$	(+)	$\sin x$
$3x^2$	(-)	$-\cos x$
$6x$	(+)	$-\sin x$
$6$	(-)	$\cos x$
$0$		$\sin x$

Again we combine the products of the functions connected by the arrows according to the operation signs above the arrows to obtain

$$\int x^3 \sin x \, dx = -x^3 \cos x + 3x^2 \sin x + 6x \cos x - 6 \sin x + C.$$

*Now try Exercise 23.*

In Exercises 21–24, use tabular integration to find the antiderivative.

21.  $\int x^4 e^{-x} dx$   
 $(-x^4 - 4x^3 - 12x^2 - 24x - 24)e^{-x} + C$

23.  $\int x^3 e^{-2x} dx$

23

23.  $\left(-\frac{x^3}{2} - \frac{3x^2}{4} - \frac{3x}{4} - \frac{3}{8}\right)e^{-2x} + C$

Handwritten work for problem 23 showing the integration process:

$$x^3 \left(-\frac{1}{2}e^{-2x}\right) - (3x^2) \left(\frac{1}{4}e^{-2x}\right)$$

$uv = \int v du$

$u$	$dv$
$x^3$	$e^{-2x}$
$3x^2$	$-\frac{1}{2}e^{-2x}$
$6x$	$+\frac{1}{4}e^{-2x}$
$6$	$-\frac{1}{8}e^{-2x}$
$0$	$+\frac{1}{16}e^{-2x}$



$$22. \int (x^2 - 5x)e^x dx$$

$$(x^2 - 7x + 7)e^x + C$$



$$u = x^2 - 5x \quad dv = e^x dx$$

$$du = 2x - 5 dx \quad v = e^x$$

$$\int (x^2 - 5x)(e^x dx) = (x^2 - 5x)(e^x) - \int (e^x)(2x - 5 dx)$$

$$u = 2x - 5 \quad dv = e^x dx$$

$$du = 2 dx \quad v = e^x$$

$$\int (2x - 5)(e^x dx) = (2x - 5)(e^x) - \int (e^x)(2 dx)$$

$$(x^2 - 7x + 7)e^x + C$$

$$= -2e^x$$

$$\begin{aligned} \int (x^2 - 5x)(e^x dx) &= (x^2 - 5x)(e^x) - (2x - 5)(e^x) + 2e^x \\ &= \left( (x^2 - 5x) - (2x - 5) + 2 \right) e^x \\ &= (x^2 - 7x + 7)e^x + C \end{aligned}$$



21.  $\int x^4 e^{-x} dx$

$(-x^4 - 4x^3 - 12x^2 - 24x - 24)e^{-x} + C$

$u$		$du$
$x^4$	$+$	$e^{-x}$
$4x^3$	$-$	$-e^{-x}$
$12x^2$	$+$	$e^{-x}$
$24x$	$-$	$-e^{-x}$
$24$	$+$	$-e^{-x}$

①

$(x^4)(-e^{-x}) - (4x^3)(e^{-x})$   
 $+ (12x^2)(-e^{-x})$   
 $- (24x)(e^{-x})$   
 $+ (24)(-e^{-x})$