

## What you'll learn about

- Linear Motion Revisited
- General Strategy
- Consumption Over Time
- Net Change from Data
- Work

## 7.1 Integral As Net Change

We actually did this  
before, back at 5.2...



...remember these?

Answers:

$$29. \int_8^{11} 87 dt = 261 \text{ miles}$$

$$30. \int_0^{60} 25 dt = 1500 \text{ gallons}$$

$$31. \int_6^{7.5} 300 dt = 450 \text{ calories}$$

In Exercises 29–32, express the desired quantity as a definite integral and evaluate the integral using Theorem 2.

29. Find the distance traveled by a train moving at 87 mph from 8:00 A.M. to 11:00 A.M.
30. Find the output from a pump producing 25 gallons per minute during the first hour of its operation.
31. Find the calories burned by a walker burning 300 calories per hour between 6:00 P.M. and 7:30 P.M.

These were easy, mostly because the integrand was a constant.  
What if it changed over the course of time?

### EXAMPLE 1 Interpreting a Velocity Function

Figure 7.1 shows the velocity

$$\frac{ds}{dt} = v(t) = t^2 - \frac{8}{(t+1)^2} \quad \frac{\text{cm}}{\text{sec}}$$


of a particle moving along a horizontal  $s$ -axis for  $0 \leq t \leq 5$ . Describe the motion.

#### SOLUTION

**Solve Graphically** The graph of  $v$  (Figure 7.1) starts with  $v(0) = -8$ , which we interpret as saying that the particle has an initial velocity of 8 cm/sec to the left. It slows to a halt at about  $t = 1.25$  sec, after which it moves to the right ( $v > 0$ ) with increasing speed, reaching a velocity of  $v(5) \approx 24.8$  cm/sec at the end. *Now try Exercise 1(a).*

SO. It'll be good for us to be savvy with our graphing calculator....

negative direction



$[0, 5]$  by  $[-10, 30]$

**Figure 7.1** The velocity function in Example 1.

## EXAMPLE 2 Finding Position from Displacement

Suppose the initial position of the particle in Example 1 is  $s(0) = 9$ . What is the particle's position at (a)  $t = 1$  sec? (b)  $t = 5$  sec?

$$\begin{aligned}\text{Displacement} &= \int_0^1 v(t) dt \\ &= \int_0^1 \left( t^2 - \frac{8}{(t+1)^2} \right) dt \\ &= \left[ \frac{t^3}{3} + \frac{8}{t+1} \right]_0^1 \\ &= \frac{1}{3} + \frac{8}{2} - 8 = -\frac{11}{3}.\end{aligned}$$

During the first second of motion, the particle moves  $11/3$  cm to the left. It starts at  $s(0) = 9$ , so its position at  $t = 1$  is

$$\text{New position} = \text{initial position} + \text{displacement} = 9 - \frac{11}{3} = \frac{16}{3}.$$



## EXAMPLE 2 Finding Position from Displacement

Suppose the initial position of the particle in Example 1 is  $s(0) = 9$ . What is the particle's position at (a)  $t = 1$  sec? (b)  $t = 5$  sec?

(b) If we model the displacement from  $t = 0$  to  $t = 5$  in the same way, we arrive at

$$\text{Displacement} = \int_0^5 v(t) dt = \left[ \frac{t^3}{3} + \frac{8}{t+1} \right]_0^5 = 35.$$

The motion has the net effect of displacing the particle 35 cm to the right of its starting point. The particle's final position is

$$\begin{aligned} \text{Final position} &= \text{initial position} + \text{displacement} \\ &= s(0) + 35 = 9 + 35 = 44. \end{aligned}$$

### EXAMPLE 3 Calculating Total Distance Traveled

Find the *total distance traveled* by the particle in Example 1.

#### SOLUTION

**Solve Analytically** We partition the time interval as in Example 2 but record every position shift as *positive* by taking absolute values. The Riemann sum approximating total distance traveled is

$$\sum |v(t_k)| \Delta t,$$

and we are led to the integral


$$\text{Total distance traveled} = \int_0^5 |v(t)| dt = \int_0^5 \left| t^2 - \frac{8}{(t+1)^2} \right| dt.$$

**Evaluate Numerically** We have

$$\text{NINT} \left( \left| t^2 - \frac{8}{(t+1)^2} \right|, t, 0, 5 \right) \approx 42.59.$$

Now try Exercise 1(c).

Why do we take the absolute value?



[0, 5] by [-10, 30]

h in


Compare the motion of the particle to the graph of  $v(t)$ ...

1.  $v(t) = 5 \cos t, \quad 0 \leq t \leq 2\pi$

$s(0) = 3$

Exercises 1–8, the function  $v(t)$  is the velocity in m/sec of a particle moving along the  $x$ -axis. Use analytic methods to do each of

b)  $\int_0^{2\pi} 5 \cos t \, dt + 3$   
 c)  $\int_0^{2\pi} |5 \cos t| \, dt$



1. (a) Right:  $0 \leq t < \pi/2, 3\pi/2 < t \leq 2\pi$   
 Left:  $\pi/2 < t < 3\pi/2$   
 Stopped:  $t = \pi/2, 3\pi/2$

(b) 0; 3                      (c) 20

2. (a) Right:  $0 < t < \pi/3$   
 Left:  $\pi/3 < t \leq \pi/2$   
 Stopped:  $t = 0, \pi/3$

(b) 2; 5                      (c) 6

7.

In Exercises 1–8, the function  $v(t)$  is the velocity in m/sec of a particle moving along the  $x$ -axis. Use analytic methods to do each of the following:

(a) Determine when the particle is moving to the right, to the left, and stopped.

(b) Find the particle's displacement for the given time interval. If  $s(0) = 3$ , what is the particle's final position?

(c) Find the total distance traveled by the particle.

1.  $v(t) = 5 \cos t$ ,  $0 \leq t \leq 2\pi$  See page 389.

2.  $v(t) = 6 \sin 3t$ ,  $0 \leq t \leq \pi/2$  See page 389.

3.  $v(t) = 49 - 9.8t$ ,  $0 \leq t \leq 10$  See page 389.

4.  $v(t) = 6t^2 - 18t + 12$ ,  $0 \leq t \leq 2$  See page 389.

5.  $v(t) = 5 \sin^2 t \cos t$ ,  $0 \leq t \leq 2\pi$  See page 389.

6.  $v(t) = \sqrt{4-t}$ ,  $0 \leq t \leq 4$  See page 389.

7.  $v(t) = e^{\sin t} \cos t$ ,  $0 \leq t \leq 2\pi$  See page 389.

8.  $v(t) = \frac{t}{1+t^2}$ ,  $0 \leq t \leq 3$  See page 389.

1. (a) Right:  $0 \leq t < \pi/2, 3\pi/2 < t \leq 2\pi$

Left:  $\pi/2 < t < 3\pi/2$

Stopped:  $t = \pi/2, 3\pi/2$

(b) 0; 3 (c) 20

2. (a) Right:  $0 < t < \pi/3$

Left:  $\pi/3 < t \leq \pi/2$

Stopped:  $t = 0, \pi/3$

(b) 2; 5 (c) 6

3. (a) Right:  $0 \leq t < 5$

Left:  $5 < t \leq 10$

Stopped:  $t = 5$

(b) 0; 3 (c) 245

4. (a) Right:  $0 \leq t < 1$

Left:  $1 < t < 2$

Stopped:  $t = 1, 2$

(b) 4; 7 (c) 6



5. (a) Right:  $0 < t < \pi/2, 3\pi/2 < t < 2\pi$

Left:  $\pi/2 < t < \pi, \pi < t < 3\pi/2$

Stopped:  $t = 0, \pi/2, \pi, 3\pi/2, 2\pi$

(b) 0; 3 (c) 20/3

2.  $v(t) = 6 \sin 3t$ ,  $0 \leq t \leq \pi/2$  See page 389.




2. (a) Right:  $0 < t < \pi/3$   
 Left:  $\pi/3 < t \leq \pi/2$   
 Stopped:  $t = 0, \pi/3$   
 (b) 2; 5

(c) 6

2) b)  $\int_0^{\pi/2} 6 \sin 3t \, dt + 3$

$= 2 + 3 = 5$

c)





In Exercises 1–8, the function  $v(t)$  is the velocity in m/sec of a particle moving along the  $x$ -axis. Use analytic methods to do each of the following:

(a) Determine when the particle is moving to the right, to the left, and stopped.

(b) Find the particle's displacement for the given time interval. If  $s(0) = 3$ , what is the particle's final position?

(c) Find the total distance traveled by the particle.

1.  $v(t) = 5 \cos t$ ,  $0 \leq t \leq 2\pi$  See page 389.

2.  $v(t) = 6 \sin 3t$ ,  $0 \leq t \leq \pi/2$  See page 389.

3.  $v(t) = 49 - 9.8t$ ,  $0 \leq t \leq 10$  See page 389.

4.  $v(t) = 6t^2 - 18t + 12$ ,  $0 \leq t \leq 2$  See page 389.

5.  $v(t) = 5 \sin^2 t \cos t$ ,  $0 \leq t \leq 2\pi$  See page 389.

6.  $v(t) = \sqrt{4 - t}$ ,  $0 \leq t \leq 4$  See page 389.

7.  $v(t) = e^{\sin t} \cos t$ ,  $0 \leq t \leq 2\pi$  See page 389.

8.  $v(t) = \frac{t}{1 + t^2}$ ,  $0 \leq t \leq 3$  See page 389.

6. (a) Right:  $0 \leq t < 4$

Left: never

Stopped:  $t = 4$

(b)  $16/3$ ;  $25/3$

(c)  $16/3$

7. (a) Right:  $0 \leq t < \pi/2$ ,  $3\pi/2 < t \leq 2\pi$

Left:  $\pi/2 < t < 3\pi/2$

Stopped:  $t = \pi/2, 3\pi/2$

(b) 0; 3

(c)  $2e - (2/e) \approx 4.7$

8. (a) Right:  $0 < t \leq 3$

Left: never

Stopped:  $t = 0$

(b)  $(\ln 10)/2 \approx 1.15$ ; 4.15

(c)  $(\ln 10)/2 \approx 1.15$

9. An automobile accelerates from rest at  $1 + 3\sqrt{t}$  mph/sec for 9 seconds.

$$v(0) = 0$$

(a) What is its velocity after 9 seconds? 63 mph

(b) How far does it travel in those 9 seconds? 344.52 feet

velocity acceleration

$$v(t) = \int a(t) dt$$

$$= \int (1 + 3\sqrt{t}) dt$$

$$= t + 2t^{3/2}$$

$$v(t) = t + 2t^{3/2}$$

total distance =  $\int_0^9 t + 2t^{3/2} dt$

$$v(t) = t + 2t^{3/2}$$

$$v(9) = 9 + 2(9)^{3/2}$$

$$9 + 2(27)$$

$$9 + 54 = 63$$

10. A particle travels with velocity

$$v(t) = (t - 2) \sin t \text{ m/sec}$$

for  $0 \leq t \leq 4$  sec.

(a) What is the particle's displacement?  $\approx -1.44952$  meters

(b) What is the total distance traveled?  $\approx 1.91411$  meters


$s(t) = \int v(t)$   $\leftarrow$  absolute value!

$$= \int_0^4 (t-2) \sin t dt$$


$$= -1.44952$$

$$\int |v(t)|$$

$$\int_0^4 |(t-2) \sin t| dt$$



In Exercises 12–16, a particle moves along the  $x$ -axis (units in cm). Its initial position at  $t = 0$  sec is  $x(0) = 15$ . The figure shows the graph of the particle's velocity  $v(t)$ . The numbers are the *areas* of the enclosed regions.



$$\textcircled{12} -4 + 5 - 24 = -23$$

$$\textcircled{13} 4 + 5 + 24 = 33$$


12. What is the particle's displacement between  $t = 0$  and  $t = c$ ?  $-23$  cm
13. What is the total distance traveled by the particle in the same time period?  $33$  cm
14. Give the positions of the particle at times  $a$ ,  $b$ , and  $c$ .  $a: 11$   $b: 16$   $c: -8$
15. Approximately where does the particle achieve its greatest positive acceleration on the interval  $[0, b]$ ?  $t = a$
16. Approximately where does the particle achieve its greatest positive acceleration on the interval  $[0, c]$ ?  $t = c$

In Exercises 17–20, the graph of the velocity of a particle moving on the  $x$ -axis is given. The particle starts at  $x = 2$  when  $t = 0$ .

(a) Find where the particle is at the end of the trip.

(b) Find the total distance traveled by the particle.

17.  $v$  (m/sec)




(a) 6 (b) 4 meters

$$\Rightarrow 4 + 2 = 6$$


$$b) = 4$$

18.




(a) 2 (b) 4 meters

19.



(a) 5 (b) 7 meters

20.



(a) -2.5 (b) 19.5 meters

**21. U.S. Oil Consumption** The rate of consumption of oil in the United States during the 1980s (in billions of barrels per year) is modeled by the function  $C = 27.08 \cdot e^{t/25}$ , where  $t$  is the number of years after January 1, 1980. Find the total consumption of oil in the United States from January 1, 1980 to January 1, 1990.  $\approx 332.965$  billion barrels

**22. Home Electricity Use** The rate at which your home consumes electricity is measured in kilowatts. If your home consumes electricity at the rate of 1 kilowatt for 1 hour, you will be charged for 1 “kilowatt-hour” of electricity. Suppose that the average consumption rate for a certain home is modeled by the function  $C(t) = 3.9 - 2.4 \sin(\pi t/12)$ , where  $C(t)$  is measured in kilowatts and  $t$  is the number of hours past midnight. Find the average daily consumption for this home, measured in kilowatt-hours.  $93.6$  kilowatt-hours

**23. Population Density** Population density measures the number of people per square mile inhabiting a given living area.

Washerton's population density, which decreases as you move away from the city center, can be approximated by the function  $10,000(2 - r)$  at a distance  $r$  miles from the city center.

(a) If the population density approaches zero at the edge of the city, what is the city's radius?  $2$  miles

(b) A thin ring around the center of the city has thickness  $\Delta r$  and radius  $r$ . If you straighten it out, it suggests a rectangular strip. Approximately what is its area?  $2\pi r\Delta r$

(c) **Writing to Learn** Explain why the population of the ring in part (b) is approximately

$$10,000(2 - r)(2\pi r) \Delta r.$$

(d) Estimate the total population of Washerton by setting up and evaluating a definite integral.  $\approx 83,776$

**24. Oil Flow** Oil flows through a cylindrical pipe of radius 3 inches, but friction from the pipe slows the flow toward the outer edge. The speed at which the oil flows at a distance  $r$  inches from the center is  $8(10 - r^2)$  inches per second.

(a) In a plane cross section of the pipe, a thin ring with thickness  $\Delta r$  at a distance  $r$  inches from the center approximates a rectangular strip when you straighten it out. What is the area of the strip (and hence the approximate area of the ring)?  $2\pi r\Delta r$

(b) Explain why we know that oil passes through this ring at approximately  $8(10 - r^2)(2\pi r)\Delta r$  cubic inches per second.

(c) Set up and evaluate a definite integral that will give the rate (in cubic inches per second) at which oil is flowing through the pipe.  $396\pi$  in<sup>3</sup>/sec or  $\approx 1244.07$  in<sup>3</sup>/sec

$$24. (b) 8(10 - r^2) \text{ in/sec} \cdot (2\pi r)\Delta r \text{ in}^2 = \text{flow in in}^3/\text{sec}$$