8-7 Study Guide and Intervention

Solving $ax^2 + bx + c = 0$

Factor $ax^2 + bx + c$ To factor a trinomial of the form $ax^2 + bx + c$, find two integers, m and p whose product is equal to ac and whose sum is equal to b. If there are no integers that satisfy these requirements, the polynomial is called a **prime polynomial**.

Example 1 Factor $2x^2 + 15x + 18$.

In this example, a=2, b=15, and c=18. You need to find two numbers that have a sum of 15 and a product of $2 \cdot 18$ or 36. Make a list of the factors of 36 and look for the pair of factors with a sum of 15.

Factors of 36	Sum of Factors
1, 36	37
2, 18	20
3, 12	15

Use the pattern $ax^2 + mx + px + c$, with a = 2, m = 3, p = 12, and c = 18.

$$2x^{2} + 15x + 18 = 2x^{2} + 3x + 12x + 18$$
$$= (2x^{2} + 3x) + (12x + 18)$$
$$= x(2x + 3) + 6(2x + 3)$$
$$= (x + 6)(2x + 3)$$

Therefore, $2x^2 + 15x + 18 = (x + 6)(2x + 3)$.

Example 2 Factor $3x^2 - 3x - 18$.

Note that the GCF of the terms $3x^2$, 3x, and 18 is 3. First factor out this GCF.

$$3x^2 - 3x - 18 = 3(x^2 - x - 6).$$

Now factor $x^2 - x - 6$. Since a = 1, find the two factors of -6 with a sum of -1.

Factors of -6	Sum of Factors
1, -6	-5
-1, 6	5
-2, 3	1
2, -3	-1

Now use the pattern (x + m)(x + p) with m = 2 and p = -3.

$$x^2 - x - 6 = (x + 2)(x - 3)$$

The complete factorization is $3x^2 - 3x - 18 = 3(x + 2)(x - 3)$.

$$3x - 3x - 10 = 3(x + 2)(x + 3)$$

Exercises

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Factor each polynomial, if possible. If the polynomial cannot be factored using integers, write *prime*.

1.
$$2x^2 - 3x - 2$$
 (2x + 1)(x - 2)

4.
$$6x^2 + 5x - 6$$
 (2x + 3)(3x - 2)

7.
$$2a^2 + 5a + 3$$
 (2a + 3)(a + 1)

10.
$$8x^2 - 4x - 24$$

2(2x - 4)(2x + 3)

13.
$$3y^2 - 6y - 24$$

3($y + 2$)($y - 4$)

16.
$$6x^2 - 7x + 18$$
 prime

2.
$$3m^2 - 8m - 3$$
 (3m + 1)(m - 3)

5.
$$3x^2 + 2x - 8$$
 (3x - 4)(x + 2)

8.
$$18y^2 + 9y - 5$$
 (6y + 5)(3y - 1)

11.
$$28p^2 + 60p - 25$$
 (2p + 5)(14p - 5)

14.
$$4x^2 + 26x - 48$$

2(x + 8)(2x - 3)

17.
$$2a^2 - 14a + 18$$

2($a^2 - 7a + 9$)

3.
$$16r^2 - 8r + 1$$
 (4r - 1)(4r - 1)

6.
$$18x^2 - 27x - 5$$
 (3x - 5)(6x + 1)

9.
$$-4t^2 + 19t - 21$$

(4t - 7)(3 - t)

12.
$$48x^2 + 22x - 15$$
 (6x + 5)(8x - 3)

15.
$$8m^2 - 44m + 48$$
 4(2m - 3)(m - 4)

18.
$$18 + 11y + 2y^2$$
 prime

Study Guide and Intervention (continued)

Solving $ax^2 + bx + c = 0$

Solve Equations by Factoring Factoring and the Zero Product Property can be used to solve some equations of the form $ax^2 + bx + c = 0$.

Example

Solve $12x^2 + 3x = 2 - 2x$. Check your solutions.

$$12x^2 + 3x = 2 - 2x$$

Original equation

$$12x^2 + 5x - 2 = 0$$

Rewrite equation so that one side equals 0.

$$(3x + 2)(4x - 1) = 0$$

Factor the left side.

$$3x + 2 = 0$$
 or $4x - 1 = 0$
 $x = -\frac{2}{3}$ $x = \frac{1}{4}$

Zero Product Property

The solution set is $\left\{-\frac{2}{3}, \frac{1}{4}\right\}$.

Since $12\left(-\frac{2}{3}\right)^2 + 3\left(-\frac{2}{3}\right) = 2 - 2\left(-\frac{2}{3}\right)$ and $12\left(\frac{1}{4}\right)^2 + 3\left(\frac{1}{4}\right) = 2 - 2\left(\frac{1}{4}\right)$, the solutions check.

Exercises

Solve each equation. Check the solutions.

$$1.8x^2 + 2x - 3 = 0$$

$$\left\{\frac{1}{2}, -\frac{3}{4}\right\}$$

4. $4x^2 = x + 3$

 $\left\{1,-\frac{3}{4}\right\}$

 $\left\{-8, \frac{5}{2}\right\}$

7. $2k^2 - 40 = -11k$

10. $12x^2 - 15 = -8x$

13. $8x^2 + 5x = 3 + 7x$

 $\left\{-\frac{3}{2}, \frac{5}{6}\right\}$

 $\left\{\frac{3}{4}, -\frac{1}{2}\right\}$

$$2. 3n^2 - 2n - 5 = 0$$

$$\left\{-1,\frac{5}{3}\right\}$$

$$5. 3x^2 - 13x = 10$$

$$\left\{-\frac{2}{3},5\right\}$$

5.
$$3x^2 - 13x = 10$$

$$\left\{-\frac{2}{3},5\right\}$$

$$8. \ 2p^2 = -21p - 40$$

$$\left\{-\frac{5}{2}, -8\right\}$$

11.
$$7a^2 = -65a - 18$$

$$\left\{-\frac{2}{7}, -9\right\}$$

14. $4a^2 - 18a + 5 = 15$

$$\left\{-\frac{1}{2},5\right\}$$

$$3. 2d^2 - 13d - 7 = 0$$

$$\left\{-\frac{1}{2},7\right\}$$

$$6.6x^2 - 11x - 10 = 0$$

$$\left\{-\frac{2}{3},\frac{5}{2}\right\}$$

$$9. -7 - 18x + 9x^2 = 0$$

$$\left\{\frac{7}{3}, -\frac{1}{3}\right\}$$

12.
$$16y^2 - 2y - 3 = 0$$

$$\left\{\frac{1}{2}, -\frac{3}{8}\right\}$$

15.
$$3b^2 - 18b = 10b - 49$$

$$\left\{\frac{7}{3}, 7\right\}$$

- **16.** The difference of the squares of two consecutive odd integers is 24. Find the integers. -5, -7 and 5, 7
- 17. GEOMETRY The length of a Charlotte, North Carolina, conservatory garden is 20 yards greater than its width. The area is 300 square yards. What are the dimensions? 30 yd by 10 yd
- **18. GEOMETRY** A rectangle with an area of 24 square inches is formed by cutting strips of equal width from a rectangular piece of paper. Find the dimensions of the new rectangle if the original rectangle measures 8 inches by 6 inches. 6 in. by 4 in.

